

# Quantitative Assessment of Fimbriae-Specific Serum and Egg Yolk Antibodies Induced in Chicken Layers by a Newly Developed Live *Salmonella* Enteritidis Vaccine and Relationship to Infection

Zeina G. Kassaify<sup>a</sup>

Ghida Banat<sup>b</sup>

Elias Baydoun<sup>c</sup>

Elie K. Barbour<sup>b</sup>

*a Nutrition and Food Sciences Department,  
Faculty of Agricultural and Food Sciences,  
American University of Beirut, Riad el Solh 1107-2020,  
PO Box: 11-0236, Beirut, Lebanon*

*b Animal Science Department,  
Faculty of Agricultural and Food Sciences,  
American University of Beirut*

*c Biology Department,  
Faculty of Arts and Sciences, American University of Beirut*

**KEY WORDS:** : Chicken layers; *Salmonella* Enteritidis vaccine; antibody; fimbriae; protection

## ABSTRACT

The increase in *Salmonella* Enteritidis (SE) human outbreaks, mostly incriminating the poultry as the source of the infection, has increased the search to control this pathogen in chickens. The purpose of this study was to quantify the SE fimbriae-specific serum and egg yolk antibodies following a controlled SE-challenge of laying hens previously immunized with a newly developed live SE vaccine versus a classical killed commercial vaccine. This work also aimed to determine the relationship between this quantitative assessment and the protection against SE

invasiveness in the livers and spleens of the experimental hens. In the procedure, laying hens were divided into four different groups. Birds in group A were given orally a newly developed live SE vaccine at 222 days of age, while birds in group D were given subcutaneously a commercial killed SE vaccine in the neck at 40 and 130 days of age. Birds of groups B and C were deprived of vaccination (control groups). Hens in groups A, B, and D were challenged orally at 237 days of age with a highly invasive SE strain, acquiring 14.1 and ~ 50 kb plasmids, at  $1.5 \times 10^{11}$  cfu/hen, while birds in group C were left without challenge. The vaccinated and challenged groups A and D and unvaccinated-challenged group B showed quantita-

tive serum antibodies-specific to *Salmonella* Enteritidis fimbriae SEF 14 and SEF 21 and egg yolk antibodies-specific to SEF 21 at 7 days post challenge. Conversely, only the live SE vaccinated-challenged group A showed quantitative egg yolk antibodies-specific to SEF 14 at 7 days post challenge. The SEF 14 and SEF 21 – specific serum antibodies decayed consistently at 14 days post challenge in groups A, B and, D. On the contrary, there was an increase in egg yolk antibodies in groups A, B and, D specific to SEF 14 and a decrease to SEF 21 at 14 days post-challenge. The highest level of SEF 14 and SEF 21 – specific egg yolk antibodies at 7 and 14 days post-challenge were recognized in group A administered the newly developed live SE vaccine. Hens of group D administered the killed SE vaccine had the highest level of SEF 14 and SEF 21 – specific serum antibodies at 7 and 14 days post-challenge. Hens in groups A and D did result in 100% protection against invasiveness of SE to livers and spleens. However, SE-challenged hens of group B, deprived of live or killed vaccine, showed a 40% and 20% invasiveness in the livers and spleens, respectively. The controlled unvaccinated and unchallenged group C didn't show any infection in livers or spleens by SE.

## INTRODUCTION

The incidence of *Salmonella* enterica serovar Enteritidis (SE) isolation associated with food illness has dramatically increased in many countries since the mid to late 1980s.<sup>1,2</sup> The U S Department of Agriculture Food Safety and Inspection Service (FSIS) data on SE in broiler chicken carcass rinses collected from 2000 through 2005 showed an annual number of isolates increase of more than four fold, and the proportion of establishments with SE-positive rinses increased nearly three fold.<sup>3</sup>

It is essential to reduce the public health risk associated with consumption of infected eggs by controlling SE infections in laying hens. Vaccination of laying hens against SE might be the most effective way to reduce egg contamination. Some live and inacti-

vated vaccines have been proven to protect against SE infection in laying hens and to reduce egg contamination.<sup>4</sup>

The fimbriae structures, located external to the cell wall of bacteria, are considered important as they frequently have been shown to mediate adhesion of SE bacterium to the host tissues.<sup>5-8</sup> The correlation between the presence of fimbriae and bacterial virulence was established in many research works.<sup>9-11</sup> Type 1 fimbriae have been shown to contribute to the intestinal stage of SE infection of chickens and to play a role in the interaction of SE with oviduct and colonization of reproductive organs.<sup>4</sup> De Buck et al (2003)<sup>12</sup> suggested that SE adhesion to the isthmal secretions was mediated by type one fimbriae and that this binding could play a role in the contamination of eggs through incorporation of the bacteria in the shell membranes.

*Salmonella* Enteritidis fimbriae 14 (SEF 14) are defined as thin fimbriae produced by the SefA gene in SE and a few other related group D serovars.<sup>13</sup> The type-1 fimbriae or SEF 21 enables the bacterium to adhere to a wide variety of eukaryotic cells in vitro.<sup>14</sup> In studies on adherence and pathogenesis of SE in mice, it was reported that SE adherence to the mouse intestinal cells involves two types of fimbriae, thus emphasizing their role in pathogenesis of infection.<sup>15</sup>

It has been reported that immunization of hens with the 14 kDa fimbrial protein results in egg-yolk antibodies specific to this protein. These antibodies were effective in inhibiting the attachment of SE to mouse intestinal epithelial cells.<sup>16</sup> The protective role of fimbriae proteins against SE infection needs to be examined in chickens.

Recently, it was shown that the early immune response in chicken to SE is not specific to any of the reported fimbriae proteins, which could explain the low immunogenicity of the fimbriae proteins leading to failure of protection in chicken against SE infection.<sup>17</sup> The greatest amount of interest in options for intervention to reduce the incidence of SE infection in chickens has been

directed toward vaccination.<sup>18</sup> The costs or impracticality of improvements in hygiene and management together with the increasing problems of antibiotic resistance suggest that vaccination in poultry against SE will become more attractive as an adjunct to existing control measures.<sup>19,20</sup>

Different preparations of classical killed vaccines for SE,<sup>14,21,22</sup> and the recent immunopotentialization of killed SE vaccine by thymulin and Zn still showed incomplete protection against infection in chicken by SE organisms.<sup>23</sup> Live attenuated vaccines against salmonellosis have been more effective in reducing mortality and shedding in challenge birds than killed vaccines.<sup>20</sup> A primary aim of developing attenuated live SE vaccines is to protect the reproductive tract and prevent the vertical transmission of this serotype. Live vaccines may invade host cells, and their efficacy may be due to their particular distribution within the body, as well as to their capability of stimulation of the cell-mediated immunity.<sup>20,24</sup>

This is the first study that quantifies serum and egg yolk antibodies specific to SEF 14 and SEF 21 fimbriae in chicken layers in response to a live *Salmonella* Enteritidis vaccine. The relationship of the quantified antibodies specific to the SE-fimbriae to protection against a controlled challenge by a highly invasive SE possessing the 14.1 and ~ 50 kb plasmids.<sup>25,26</sup>

## **MATERIALS AND METHODS**

### **Chicken Layers**

A total of 20 brown feathered Babcock layers of 222 days of age were provided by the Agricultural Research and Education Center (AREC) Bekaa, Lebanon. The hens were divided into four groups (5 hens/group) each group was put in a separate isolated room. All groups of hens were fed similarly according to NRC, 1994 standard requirement.<sup>27</sup> Cloacal swabs collected from all the hens and the laboratory culture results confirmed the hens as free from *Salmonella* infections.

### **Newly Developed SE Vaccine**

The SE isolate used was chosen according to its prevalence; it had the most common plasmid profile among SE isolates recovered from eleven poultry broiler flock outbreaks.<sup>25</sup> The isolate had two plasmids located at different base pair positions, namely, 14.1 and ~50 kb. The attenuation of the field highly invasive SE strain acquiring 14.1 and ~50 kb plasmids was performed in a low concentration of formalin equivalent to 0.38x10<sup>-3</sup>%. The contact time between the SE cells and highly diluted formalin, at room temperature, was 15 hours, resulting in a viable count of the attenuated SE cells equivalent to 2.4x10<sup>8</sup> cfu/ml.

### **Vaccination**

The four groups of hens were labeled as A, B, C, and D, and each group received a different treatment. The hens in group A received 1 ml of the live SE vaccine orally (2.4x10<sup>8</sup> cfu/ml) followed by 3 ml of saline to wash in the vaccine. The hens in groups B and C were left without vaccination; however, hens of group D received a classical killed commercial SE vaccine (Hipra, Spain). The killed SE vaccine was administered to the hens subcutaneously through the neck at a level of 0.5 ml/hen and delivered twice, at 40 and 130 days of age, according to the manufacturer's instructions.

### **Challenge Inoculum**

A highly virulent and invasive SE isolate was used for the preparation of the challenge inoculum. The involvement of this SE isolate, acquiring plasmids 14.1 and ~ 50 Kb, in severe outbreaks of poultry, has been reported previously to the World Animal Health Organization by the authors of this work.<sup>25</sup> Two passages in vivo were performed to the isolate in order to raise its virulence. In brief, 24 hr-culture of SE colonies that were heavily grown on a 15 cm-diameter Brilliant Green Agar (BGA) (HIMedia Laboratories, Mumbai, India) plate were scraped from 1/3rd of the plate area. The cell mass was then reconstituted in 1 ml of sterile saline and 0.5 ml was delivered intravenously in the brachial vein of a 2-months-old white Pop Quail.

The invasive SE was recovered from the spleen by culturing on BGA. The recovered colony was subcultured by heavy seeding on another BGA plate that was incubated for 24 hrs at 37°C. One third of the seeded colonies were scraped from the BGA plate and reconstituted in 0.5 ml sterile saline. Another 2-month-old white Pop Quail was intravenously inoculated with 0.5 ml of the SE suspension in its brachial vein. The SE was recovered from the spleen of the infected white Pop Quail on a BGA plate that was incubated for 24 hrs at 37°C. All the SE recovered colonies were scraped from the BGA plate and reconstituted in 100 ml of sterile Tryptose Phosphate Broth (Himedia Laboratories, Mumbai, India). The SE count in the prepared challenge was  $1.5 \times 10^{11}$  cfu/ml.

Each hen in groups A, B, and D received orally 1 ml of the SE challenge ( $1.5 \times 10^{11}$  cfu/ml/hen) at 237 days of age, followed by 3 ml of sterile saline to wash in the challenge, while group C remained unvaccinated and unchallenged as a negative control group.

#### **Immunoblott Sera and Egg Yolk**

Blood samples were collected from the brachial vein of all the hens in the four different groups at 7 and 14 days post-challenge (244 and 251 days of age). The five serum samples collected from the five hens of the same age belonging to each group were pooled in equal portions. Each pooled sera was stored at -20°C for use in Western Immunoblotting.

The eggs were collected at 7 and 14 days post-SE challenge, transported in refrigerated containers, cleaned with potable water using a detergent and a disinfectant, and then stored in a refrigerator in cleaned marked bags until the experimental testing was performed.

Before analysis, each collected egg was disinfected with 70% alcohol and a sterile cotton swab. A hole was then made in the eggshell at the aircell side using a sterile forceps and scissors. After removing a piece of the vitelline membrane, a 1 ml of the

yolk was obtained using a clean disposable pipette and mixed with 1 ml of phosphate buffered saline (PBS) and vortexed for 1 minute. The above mixture was centrifuged at 2,500 rpm for 30 minutes using an IEC centrifuge (HN-SII, Needham, HTS., MA, USA). The supernatant was collected. The supernatants of the egg yolk samples that were collected from hens of same age in each group were pooled in equal portions. Each pooled egg yolk sample was stored at -20°C to be used in Western Immunoblotting.

#### **Isolation of SE Fimbriae Fourteen and Twenty one KDa**

*Salmonella* Enteritidis was grown statically in 2 liters of Colonization Factor Antigen medium (Difco, Becton Dickinson, Sparks, MD, USA) at 37°C for 60 hrs. The SE culture was harvested by centrifugation at 3,000 rpm (180xg) for 10 minutes using a GLC-2B centrifuge (Du Pont Instruments, Sorvall®, Newtown, CT, USA), then suspended in 120 ml of 0.15 M ethanolamine buffer, pH 10.5.13 Fimbriae were separated from the SE cells at room temperature (25°C) by shearing them in a blender (mixer Blender 3, Mammonlex, Model 242, Taiwan) for three cycles (1 minute/cycle). The cells and cellular debris were removed by centrifugation at 12,000xg for 15 minutes at 4°C using RC2-B Automatic Refrigerated Super Speed Centrifuge by Sorvall®.

The supernatant (Fraction 1) was centrifuged at 100,000xg for 1 hr at 4°C using OTD 65B ultracentrifuge by Sorvall® in order to remove the membrane vesicles. The result is a clarified supernatant (Fraction 2), which was dialyzed overnight in the fridge against 10 mM Tris HCl (pH 7.5) containing 0.2% SDS in a dialysis bag (Spectrum Lab, Rancho Dominguez, CA, USA) of molecular weight pores cutoff between 6-8 KDa, in order to precipitate the SEF 14. The SEF 14 fimbriae were pelleted by centrifugation at 15,000xg for 15 minutes at 4°C using RC-5B centrifuge by Sorvall®, thus, separating the precipitated SEF 14 from the SEF 21, which remained in the supernatant

as fractions<sup>3</sup>. The SEF 14 pellets were then reconstituted in 5 ml sterile saline. On the other hand, fraction 3 containing the SEF 21 fimbriae (75ml) was concentrated to approximately 25 ml by dialysis against 30% polyethylene glycol 20,000 (Sigma, St. Louis, MO, USA) for around 2 hours followed by precipitation of SEF 21 with 37.5 ml of ice-cold acetone. The precipitated SEF 21 was recovered by centrifugation at 15,000xg for 20 minutes at 4°C. The pellet (fraction 4), containing the SEF 21, was suspended in 4 ml of Laemmli buffer and boiled for 5 minutes to solubilize the constituents.<sup>13</sup>

### **SDS-PAGE of Isolated Fimbriae**

The banding of the SEF 14 and 21 KDa fimbriae was done by Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE), using the discontinuous buffer system.<sup>28</sup> The weight of protein of SEF 14 applied in 20 µl volume per lane was 1.75 µg, while that of SEF 21 was 3.5 µg. The molecular weight marker was diluted 1:20 with SDS reducing buffer and applied on gel in 10µl volume. A 12% separating gel was allowed to polymerize for 45 minutes in a mini-protean II electrophoresis cell (Bio-Rad Lab., Richmond, CA, USA). The electric current in the gel was run at 60 mA for 45 minutes.

### **Serum and Egg Yolk Antibodies to Fimbriae**

The detection of chicken serum and egg yolk antibodies to SEF 14 and SEF 21 was performed by Western immunoblotting.<sup>29</sup> Briefly, the fimbriae of SE resolved on SDS-PAGE gels were electrophoretically transferred onto a nitrocellulose membrane (NCM) (Bio-Rad Lab, Richmond, CA, USA). The electrophoretic transfer was performed in a transblot cell (Bio-Rad Lab., Richmond, CA, USA) for 1 hr. at 0.25 A and 100 V. Blocking of the active sites was performed by immersion of the NCM in 5% gelatin-Tris-Buffer Saline for 2 hrs at 37°C. The pooled sera and egg yolk samples collected from each group of hens at a specific time (7 and 14 days post SE challenge) were diluted to 1:250 and 1:50, respec-

tively, using 1% gelatin-Tris-Tween-Buffer Saline (TTBS). The SEF-14 and SEF-21 lanes were individually reacted to each pooled sera and egg yolk samples for 10 hrs at 37°C. A goat anti-chicken IgG (H+L) peroxidase conjugate (Sigma, St. Louis, MO, USA) was diluted to 1:1000 in 1% gelatin-TTBS and then added to NCM and incubated at 37°C for 1 hr. The substrate used was 3,3'-DAB peroxidase (Sigma, St. Louis, MO, USA), added to NCM for 30 minutes at 37°C. The NCM containing the formed brown colored bands was rinsed with distilled water and dried over a filter paper. The dried NCM-bands were scanned using Scanjet 6300C (Hewlett Packard, USA) with setting at high Sharpen level, and output resolution of 300.

### **Quantitative**

#### *Assessment of Antibodies*

The antibodies-specific to SEF 14 and SEF 21 fimbriae as colorimetrically formed on NCM were quantitatively measured by reading the mean absorbance (intensity of color) of the scanned bands using a new computerized program developed by the National Institute of Health, USA namely the NIH Image 1.62 program. This program is available on the Internet at <http://rsb.info.nih.gov/nih-image> powered by executor for windows, which is available also on the Internet at <http://www.ardi.com>. The mean intensity values obtained for antibodies specific to the banded fimbriae of SE in groups A, B, C, and D were subtracted from the background mean intensity of the control group "C" at each specific age.

#### *Protection*

To assess the invasiveness of the highly virulent field SE isolate acquiring 14.1 and ~50 kb plasmids in the laying hens that were orally challenged, SE was attempted to be recovered from the livers and spleens of those hens. All the laying hens in the four different groups were sacrificed by cervical dislocation at 14 days post SE challenge (251 days of age). An area of 0.75x1.0 cm<sup>2</sup> was cut aseptically from each organ (livers and spleens) and cultured in 5 ml of sterile

**Table 1.** Nature of treatments in the four groups of hens

Hen group	Nature of vaccine	Challenged <sup>a</sup>
A	Live <sup>b</sup>	Yes
B	NA <sup>c</sup>	Yes
C	NA	No
D	Killed <sup>d</sup>	yes

a Each challenged layer received orally  $1.5 \times 10^{11}$  cfu/ml/hen of an invasive SE strain acquiring 14.1 and ~50 Kb plasmids, at 237 days of age (14 days post live SE vaccination in hens of group A).

b A viable count of  $2.4 \times 10^8$  cfu/ml of live attenuated SE vaccine-cells per hen was administered orally at 222 days of age.

c NA = not applicable.

d Group D hens were each given a killed commercial SE vaccine (Hipra, Spain). The vaccine was delivered in 0.5 ml/hen, subcutaneously in the neck, at 40 and 130 days of age.

**Table 2.** Quantitative assessment<sup>a</sup> of chicken serum and egg yolk antibodies specific to SEF 14 in response to challenge

Hen Group <sup>b</sup>	Mean intensity of antibody response <sup>c</sup> to banded SEF 14 (Days post SE challenge / Age)			
	Serum		Egg Yolk	
	7 / 244	14 / 251	7 / 244	14 / 251
A	+12.2	-11.9	+0.3	+13.4
B	+58.2	+10.1	-9.3	-3.9
C	0.0	0.0	0.0	0.0
D	+126.5	+18.4	-5.3	+12.9

a Assessed by scanning of bands found on Nitrocellulose membrane followed by reading intensities of the bands using NIH Image 1.62 computer program

b Hen groups or treatments are defined in Table 1.

c The means of intensity of SEF 14 bands reacted to serum of hens in groups A, B, C and D is subtracted from the background mean obtained by hens of the control group "C" at each specific age.

enrichment Selenite Broth. The Selenite cultures were incubated at 41.5°C for 20-24 hrs. Individual subcultures from selenite to Brilliant Green Agar (BGA) plates were performed. The BGA culture was incubated at 37°C for 24 hrs. Suspected *Salmonella* colonies on BGA were tested biochemically for conformation of identity using H<sub>2</sub>S gas production and glucose fermentation in Triple Sugar Iron (TSI) agar (HIMedia Laboratories, Mumbai, India) and the lack of urease production in Urea Agar Base medium.<sup>30</sup>

## RESULTS

Table 2 shows the quantified chicken serum and egg yolk antibodies specific to SEF 14 fimbriae at different times post SE challenge. An apparent consistent decay in the SEF 14 serum antibodies and an increase in

the SEF 14 egg yolk antibodies in groups A, B, and D was noticed at 14 days post challenge in comparison to that at 7 days post challenge. The hens in group D that were administered two doses of killed SE bacterin showed the highest level of serum antibodies to SEF 14 at 7 and 14 days post challenge in comparison to hens of group A (administered a live vaccine and a challenge) and hens of group B (unvaccinated and challenged). On the other hand, the hens of group A showed the highest level of egg yolk antibodies to SEF 14 at 7 and 14 days post challenge in comparison to hens in groups B and D.

Serum antibodies to SEF 14 were detected earlier after challenge and in higher quantities in the hens of groups A, B, and D

**Table 3.** Quantitative assessment of chicken serum and egg yolk antibodies specific to SEF 21 in response to challenge

Hen Group <sup>b</sup>	Mean intensity of antibody response <sup>c</sup> to banded SEF 21 (Days post SE challenge / Age)			
	Serum		Egg Yolk	
	7 / 244	14 / 251	7 / 244	14 / 251
A	+23.2	+16.7	+18.6	+2.3
B	+15.5	+10.7	+6.4	-0.1
C	0.0	0.0	0.0	0.0
D	+66.2	+37.2	+3.49	+1.3

a Assessed by scanning of bands found on Nitrocellulose membrane followed by reading intensities of the bands using NIH Image 1.62 computer program.

b Hen groups or treatments are defined in Table 1.

c The means of intensity of SEF 21 bands reacted to serum of hens in groups A, B, C and D is subtracted from the background mean obtained by hens of the control group "C" at each specific age.

**Table 4.** Protection against SE invasiveness in visceral organs at 2 weeks (251 days of age) post challenge of the different groups

Hen group <sup>a</sup>	% (frequency <sup>b</sup> ) of layers with SE infection in different organs	
	Liver	Spleen
A	0 (0/5)	0 (0/5)
B	40 (2/5)	20 (1/5)
C	0 (0/5)	0 (0/5)
D	0 (0/5)	0 (0/5)

a The nature of the treatment in each of the four groups of layers is defined in Table 1.

b Frequency is the number of layers with SE recovery from the specific organ divided by the number of layers in a treatment.

compared to that of their respective egg yolk antibodies. Moreover, only the hens of group A had SEF 14 egg yolk antibodies detected in small amounts at 7 days post challenge. In addition, 14 days post challenge, SEF 14 egg yolk antibodies in group A were in higher quantities than those detected in the sera.

Table 3 shows the quantified chicken serum and egg yolk antibodies specific to SEF 21 at different times following the SE challenge. The serum antibodies specific to SEF 21 decayed consistently in groups A, B, and D with time post challenge (Table 3), similar to the consistent decay with time shown for serum antibodies specific to SEF 14 (Table 2). However, the yolk antibodies to SEF 21 raised with time post challenge (Table 3), while the yolk antibodies to SEF

14 dropped with time (Table 2).

The hens in group D that were given two doses of a killed SE bacterin, and when challenged, showed the highest level of serum antibodies to SEF 21 at 7 and 14 days post challenge in comparison to the hens in group A that were given the live vaccine and challenged and the hens in group B that were unvaccinated and challenged.

On the other hand, the hens of group A showed the highest level of egg yolk antibodies to SEF 21 at 7 and 14 days post challenge in comparison to the hens in groups B and D.

The SEF 21 specific antibodies were detected in both the serum and egg yolk of hens in groups A, B, and D as early as 7 days post challenge (Table 3). However,

higher quantities of antibodies were found in the serum rather than in their respective egg yolks.

The protection effect against SE invasiveness into the visceral organs (livers and spleens) at 14 days post challenge is shown in Table 4. The live vaccine given to hens of group A, and the killed vaccine given to hens of group D did result in 100% protection against invasiveness of SE to livers and spleens. However, the hens in group B, deprived of live or killed vaccine, showed a 40% and 20% invasiveness in the livers and spleens, respectively. These hens of the control group C were free of *Salmonella* infections in their livers and spleens, a reflection of the animal room environment standard, indicating a proper biosecurity and compliance to isolation requirements.

## DISCUSSION

The quantitative assessment of the serum antibodies specific to SEF 14 (Table 2) in the various vaccinated hens reveals the ability of both the newly developed live SE vaccine and the commercial killed SE vaccine to induce enough protective SEF 14 antibodies at 7 days post challenge as evidenced in the hens of groups A and D respectively. This effect was maintained in spite of the neutralization by the invading challenging organism. Such a finding is in agreement with prior research on the ability of vaccines to maintain immunity during neutralization by the invading organisms.<sup>31,32</sup>

On the other hand, the early yolk antibody response at 7 days post challenge to SEF 14 was only evident in eggs of group A that were given the live SE vaccine. This finding disagrees with other previous observations,<sup>33-34</sup> in which early detection of egg yolk antibodies was absent. This early immune response to SEF 14 may be due to a difference in the nature of the newly developed vaccine and/or the genotype of the experimental hens.<sup>35-36</sup>

The consistent decay of the serum antibodies specific to SEF 14 by time (14 days versus 7 days post challenge) in groups A, B, and D may be contributed to the short

half life of chicken antibodies and/or due to the neutralization effect by the invading SE organism.<sup>37</sup> On the other hand, the increase in the SEF 14 specific egg yolk antibodies by time, in groups A, B, and D, may perhaps reflect the time necessary for antibodies to transit from serum to the egg yolk during the egg formation.<sup>38</sup>

The two groups of hens, B and D, that showed the highest levels of SEF 14 specific serum antibodies at 7 days post challenge were able to keep these highest levels at 14 days post challenge. The killed vaccine administered to hens of group D was able to induce after SE challenge a higher antibody level specific to SEF 14 than the serum antibody levels obtained by group A hens that received the live attenuated SE vaccine (Table 2). It has been noted in previous studies the inability of attenuated live enteric vaccines to induce a significant humoral response than that caused by a killed vaccine.<sup>20,39</sup> However, the hens in group A that were given the live SE vaccine were able to keep a much higher level of egg yolk antibodies at 14 days post challenge in comparison to that of groups B and D. This finding is supported by some earlier studies, showing stronger and long lasting local immunity induction against *Salmonella* infection in layers by live attenuated vaccines when compared with killed bacterins.<sup>21,40-43</sup>

The serum and egg yolk antibody response to SEF 21 was evident in groups A, B, and D at 7 days post challenge (Table 3). This result could indicate a higher immunogenicity of the 21 KDa compared to the 14 KDa fimbriae. In a previous study, it was indicated that as the molecular weight of the polypeptide antigen increases its immunogenicity increases accordingly.<sup>17</sup> The SEF 21 specific serum and egg yolk antibodies in the hens of group B that were not vaccinated, but challenged, were lower than those obtained in the vaccinated and challenged groups (A and D) at all times. This significantly indicate the importance of vaccination (sensitization) that induces a secondary immune response post challenge.<sup>37,40,41,43</sup>

Hens in group D given the killed SE bacterin had the highest level of serum antibodies to SEF 21 in comparison to the other groups of hens. This is in agreement with previous reports comparing the killed to live vaccines in induction of humoral immunities.<sup>20</sup> Hens in group A, given the newly developed live vaccine, had the highest level of egg yolk antibodies to SEF 21 when compared with the other groups. This observation is in agreement with previous findings in which the live vaccines induced better local immunities than the killed bacterins.<sup>14,21,40-43</sup>

The above results confirm the ability of both the newly developed live SE vaccine and the commercial killed SE vaccine to maintain an adequate SEF 14 and SEF 21-serum antibodies at the critical period post challenge (Tables 2 and 3). This may have helped in the neutralization of the invasive SE organisms that were used in challenging the hens. Thus, leading to clearance of infection from the livers and spleens of hens in groups A and D (Table 4). It was previously noted that the presence of protective antibodies could intercept with infection and invasion of microorganisms in the host.<sup>21,22</sup> Hens in group B that were deprived of vaccination were able to induce antibodies against SEF 14 and SEF 21 (Tables 2 and 3) as a result of the infection caused by the challenging organisms. However, this infection titer was not able to protect against invasiveness of SE organisms in this group of hens (Table 4). A reason for this could be the absence of memory cells to SE due to deprivation from vaccination,<sup>44,45</sup> more specifically, the absence of sensitized T-helper and T-delayed hypersensitivity cells that are helpful in cell-mediated immune responses against intracellular infections, such as that caused by SE organisms.<sup>20</sup>

In conclusion, both the newly developed live vaccine and the commercial classical killed SE vaccine were able to maintain SEF-antibodies during the critical period of challenge. This contributed to the protectiveness against the invasion of visceral

organs by the highly virulent SE used in the challenge.

## ACKNOWLEDGEMENTS

The authors thank the University Research Board at the American University of Beirut for its interest and support of this project.

## REFERENCES

- Hogue A, White P, Guard-Petter J, Schlosser W, Gast R, Ebel E, Farrar J, Gomez T, Madden J, Madison M, McNamara AM, Morales R, Parham D, Sparling P, Sutherland W and Swerdlow D: Epidemiology and control of egg-associated *Salmonella* Enteritidis in the United States of America. *Rev Sci Tech Off Int Epiz* 1997;16:542-553
- Withanage G.K, Sasai K., Fukata T, Miyamoto T, Lillehoj HS, Baba E: Increased lymphocyte subpopulations and macrophages in the ovaries and oviducts of laying hens infected with *Salmonella enterica* serovar Enteritidis. *Avian Pathol* 2003;32:583-590
- Altekruse SF, Bauer N, Chanlongbutra A, DeSagun R, Naugle A, Schlosser W, Umholtz R, PWhite R: *Salmonella* Enteritidis in Broiler Chickens, United States, 2000-2005. *Emerg Infect Dis* 2006;12:1848-1852
- De Buck J, Van Immerseel F, Haesebrouck F, Ducatelle R: Protection of laying hens against *Salmonella* Enteritidis by immunization with type 1 fimbriae. *Vet Microbiol* 2005;105:93-101
- Feurtrier J, Kay WW, Trust TJ: Cloning and expression of a *Salmonella* enteritidis fimbria gene in *Escherichia coli*. *J Bacteriol* 1988; 170:4216-22.
- Lindquist BL, Libenthal E, Lee PC, Stinson MW, Merrick JM: Adherence of *Salmonella typhimurium* to small-intestinal enterocytes of the rat. *Infect Immun.* 1987; 55:3044-50.
- Muller KH, Collinson SK, Trust TJ, Kay WW: Type 1 fimbriae of *Salmonella enteritidis*. *J Bacteriol.* 1991; 173: 4765-72.
- Parkkinen J, Ristimake A, Westerlund B: Binding of *Escherichia coli* S fimbriae to cultured human endothelial cells. *Infect Immun.* 1989; 57:2256-9.
- Ihawi T, Abel Y, Nakao M, Imada A, Tsuchiya K: Role of type 1 fimbriae in the pathogenesis of ascending urinary tract infection induced by *Escherichia coli* in mice. *Infect Immun.* 1983; 39:1307-15.
- McCormick BA, Franklin DP, Laux DC, Cohen PS: Type 1 pilli are not necessary for colonization of the streptomycin-treated mouse large intestine by type 1 - pilliated *Escherichia coli* F-18 and *E. coli* K-12. *Infect Immun.* 1989; 57:3022-9.
- Reid G, Sobel JD: Bacterial adherence in the pathogenesis of urinary tract infection: a review. *Rev Infect Dis.* 1987; 9:470-87.
- De Buck J, Van Immerseel F, Meulemans G, Haesebrouck F and Ducatelle O: Adhesion of *Salmonella enterica* serovar Enteritidis isolates to chicken isthmal glandular secretions. *Vet Microbiol* 2003; 93:223-233.

13. Feutrier J, Kay WW, Trust TJ: Purification and characterization of fimbriae from *Salmonella* enteritidis. *J Bacteriol.* 1986; 168:221-7.
14. Barrow PA: Experimental infection of chicken with *Salmonella* enteritidis. *Avian Pathol.* 1991; 20:145-53.
15. Aslanzadeh J, Paulissen LJ: Role of type 1 and type 3 fimbriae on the adherence and pathogenesis of *Salmonella* enteritidis in mice. *Microbiol Immunol.* 1992; 36:351-9.
16. Peralta RC, Yokoyama H, Ikemori Y, Kuroki M, Kodoma Y: Passive immunisation against experimental salmonellosis in mice by orally administered hen egg-yolk antibodies specific for 14-kD fimbriae of *Salmonella* enteritidis. *J Med Microbiol.* 1994; 41: 29-35.
17. Barbour EK, El Jurdi LH, Faroon OM, Daghir NJ, Bouljihad M: Chronological recognition by chicken of antigenic polypeptides in *Salmonella* Enteritidis with different plasmid profiles: relationship to infection rate. *J Vet Med Sci.* 2000; 62(6):565-70.
18. Gast RK, Bear CW: Research to understand and control *Salmonella* Enteritidis in chickens and eggs. *Poult Sci* 1993;72:1157-1163.
19. Zhang-Barber L, Turner AK, Barrow PA: Vaccination for control of *Salmonella* in poultry. *Vaccine* 1999; 17: 2538–2545.
20. Nagaraja KV, Rajashekara G: Vaccination against *Salmonella* enterica serovar Enteritidis infection: dilemma and realities. In: Saeed AM, Gast RK, Potter ME, Wall, PG, editors. *Salmonella* Enterica serovar Enteritidis in humans and animals-epidemiology, pathogenesis, and control. Ames: Iowa State University Press, Ames, 1999. p. 397-404.
21. Barbour EK, Frerichs WM, Nabbut NH, Poss PE, Brinton MK: Evaluation of bacterins containing three predominant phage types of *Salmonella* enteritidis for prevention of infection in egg-laying chickens. *Am J Vet Res.* 1993; 54:1306-9.
22. Gast RK, Stone HD, Holt, PS: Evaluation of the efficacy of oil-emulsion bacterins for reducing fecal shedding of *Salmonella* Enteritidis in laying hens. *Avian Dis.* 1993; 37:1085-91.
23. Barbour EK, Bejjani NE, Daghir NJ, Faroon OM, Bouljihad M, Spasojevic R: Induction of early immunopotentiality to fimbriae of *Salmonella* Enteritidis (SE) by administering thymulin and zinc to SE-vaccinated chicken breeders: relationship to protection. *J Vet Med Sci.* 2000; 62(11):1139-43.
24. De Buck J, Van Immerseel F, Haesebrouck F, Ducatelle R: Colonization of the chicken reproductive tract and egg contamination by *Salmonella*. *J Appl Microbiol* 2004; 97: 233–245
25. Barbour EK, El Jurdi LH, Talhouk R, Qatanani M, Eid A, Sakr W, Bouljihad M, Spasojevic R: Emergence of *Salmonella* Enteritidis outbreaks in broiler chickens of Lebanon: epidemiologic markers and competitive exclusion control. *Sci Tech Rev J* (World Anim Hlth Org). 1999; 18:710-8.
26. Chart H, Threlfall EJ, Row B: Virulence of *Salmonella* Enteritidis phage type 4 is related to the possession of a 38 Mda plasmid. *Fed Eur Microbiol Soc Microbiol Lett.* 1989; 58:229-304.
27. NRC (National Research Council): Nutrient requirements of poultry. 8th revised ed., Washington, D.C.: National Academy Press, 1984. p. 4-15.
28. Laemmli UK: Cleavage of structural protein during the assembly of the head of bacteriophage T4. *Nature* (Lond). 1970; 227:680-5.
29. Towbin H, Stachelin T, Gordon J: Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA.* 1979; 76:4350-4.
30. American Association of Avian Pathologists: Salmonellosis. In: Swayne DE, Glisson JR, Jackwood MW, Pearson JE, Reed WM, editors. A laboratory manual for the isolation and identification of avian pathogens, 4th ed. Tallahassee, Florida: Rose Printing: Tallahassee, Florida, 1998. p. 4-13.
31. Desiderio JV, Campbell SG: Immunization against experimental murine Salmonellosis with liposome-associated O-antigen. *Infect Immun.* 1985; 48:658-63.
32. Knivett VA, Tucker JF: The evaluation of a live *Salmonella* vaccine in mice and chickens. *J Hyg.* 1991; 69:233-45.
33. Holt PS, Stone HD, Gast RK, Greene CR: Application of the agar gel precipitin test to detect antibodies to *Salmonella* enterica serovar Enteritidis in serum and egg yolks from infected hens. *Poult Sci.* 2000; 79: 1246-51.
34. Sunwoo HH, Nakano T, Dixon WT, Sim JS: Immune responses in chickens against lipopolysaccharide of *Escherichia coli* and *Salmonella typhimurium*. *Poult Sci.* 1996; 75: 342-5.
35. Gast RK: Paratyphoid infections. In: Calnek BW, Barnes HJ, Beard CW, Dougald LR, Saif YM, editors. *Diseases of poultry*, 10th Edition. Ames: Iowa State University Press, Ames, 1997. p. 97-121.
36. Gast RK: Applying Experimental Infection Models to Understand the Pathogenesis, Detection, and Control of *Salmonella* enterica Serovar Enteritidis in Poultry. In: Saeed AM, Gast RK, Potter ME, Wall PG, editors. *Salmonella* Enterica Serovar Enteritidis in Humans and Animals – Epidemiology, Pathogenesis, and Control. Ames: Iowa State University Press, Ames, 1999. p. 233-43.
37. Barbour EK, Hamadeh SK, Abi Ghanem D, Haddad JJ, Safieh-Garabedian B: Humoral and cell-mediated immunopotentiality in vaccinated chicken layers by thymic hormones and zinc. *Vaccine.* 1998; 16:1650-5.
38. Brambell FWR: The transmission of passive immunity from mother to young. North-Holland Publishing Company: Amsterdam, 1970. p. 23-8.
39. Arnon R, Shapria M, Jacob CO: Synthetic vaccine. *J Immunol Methods.* 1989; 61:261-73.
40. Barrow PA, Lovell MA, Berchieri A: Immunization of laying hens against *Salmonella* Enteritidis with live attenuated vaccines. *Avian Pathol.* 1990; 126: 241-2.
41. Hassan JO, Curtiss III R: Control of colonization by virulent *Salmonella typhimurium* by oral immunization of chickens with avirulent cya- crp- *Salmonella typhimurium*. *Res Microbiol.* 1990;

- 141:839-50.
42. Hassan JO, Curtiss III R: Efficacy of a live avirulent *Salmonella* typhimurium vaccine in preventing colonization and invasion of laying hens by *Salmonella* typhimurium and *Salmonella* Enteritidis. *Avian Dis.* 1997; 41: 783-91.
  43. Vielitz E, Conard EC, Vob M, Lohren U, Bachmeierand J, Hahn I: Immunization against *Salmonella*-infection using live and inactivated vaccine preparations. In: World Poultry Congress, Amsterdam, 1992. p. 435-8.
  44. Barrow PA, Lovell MA, Berchieri A: The use of two live attenuated vaccines to immunize egg-laying hens against *Salmonella* enteritidis phage type 4. *Avian Pathol.* 1991; 20:681-92.
  45. Finlay BB, Falkow S: Common themes in microbial pathogenicity. *Microbiol Rev.* 1989; 53:210-30.